

# Consistency of Functional Learning Methods Based on Derivatives

Fabrice Rossi<sup>a</sup>, Nathalie Villa-Vialaneix<sup>b,c,\*</sup>

<sup>a</sup>*Télécom ParisTech, LTCI - UMR CNRS 5141, France*

<sup>b</sup>*IUT de Perpignan (Dpt STID, Carcassonne), Université de Perpignan Via Domitia, France*

<sup>c</sup>*Institut de Mathématiques de Toulouse, Université de Toulouse, France*

---

## Abstract

In some real world applications, such as spectrometry, functional models achieve better predictive performances if they work on the derivatives of order  $m$  of their inputs rather than on the original functions. As a consequence, the use of derivatives is a common practice in functional data analysis, despite a lack of theoretical guarantees on the asymptotically achievable performances of a derivative based model. In this paper, we show that a smoothing spline approach can be used to preprocess multivariate observations obtained by sampling functions on a discrete and finite sampling grid in a way that leads to a consistent scheme on the original infinite dimensional functional problem. This work extends Mas and Pumo (2009) to nonparametric approaches and incomplete knowledge. To be more precise, the paper tackles two difficulties in a nonparametric framework: the information loss due to the use of the derivatives instead of the original functions and the information loss due to the fact that the functions are observed through a discrete sampling and are thus also imperfectly known: the use of a smoothing spline based approach solves these two problems. Finally, the proposed approach is tested on two real world datasets and the approach is experimentally proven to be a good solution in the case of noisy functional predictors.

*Keywords:* Functional Data Analysis, Consistency, Statistical learning, Derivatives, SVM, Smoothing splines, RKHS, Kernel

---

\*Corresponding author.

*Email addresses:* [Fabrice.Rossi@telecom-paristech.fr](mailto:Fabrice.Rossi@telecom-paristech.fr) (Fabrice Rossi),  
[nathalie.villa@math.univ-toulouse.fr](mailto:nathalie.villa@math.univ-toulouse.fr) (Nathalie Villa-Vialaneix)

*Preprint submitted to Pattern Recognition Letters*

*May 1, 2011*

---

## 1. Introduction

As the measurement techniques are developing, more and more data are high dimensional vectors generated by measuring a continuous process on a discrete sampling grid. Many examples of this type of data can be found in real world applications, in various fields such as spectrometry, voice recognition, time series analysis, etc.

Data of this type should not be handled in the same way as standard multivariate observations but rather analysed as *functional* data: each observation is a function coming from an input space with infinite dimension, sampled on a high resolution sampling grid. This leads to a large number of variables, generally more than the number of observations. Moreover, functional data are frequently smooth and generate highly correlated variables as a consequence. Applied to the obtained high dimensional vectors, classical statistical methods (e.g., linear regression, factor analysis) often lead to ill-posed problems, especially when a covariance matrix has to be inverted (this is the case, e.g., in linear regression, in discriminant analysis and also in sliced inverse regression). Indeed, the number of observed values for each function is generally larger than the number of functions itself and these values are often strongly correlated. As a consequence, when these data are considered as multidimensional vectors, the covariance matrix is ill-conditioned and leads to unstable and unaccurate solutions in models where its inverse is required. Thus, these methods cannot be directly used. During past years, several methods have been adapted to that particular context and grouped under the generic name of Functional Data Analysis (FDA) methods. Seminal works focused on linear methods such as factorial analysis (Deville (1974); Dauxois and Pousse (1976); Besse and Ramsay (1986); James et al. (2000), among others) and linear models Ramsay and Dalzell (1991); Cardot et al. (1999); James and Hastie (2001); a comprehensive presentation of linear FDA methods is given in Ramsay and Silverman (1997, 2002). More recently, nonlinear functional models have been extensively developed and include generalized linear models James (2002); James and Silverman (2005), kernel nonparametric regression Ferraty and Vieu (2006), Functional Inverse Regression Ferré and Yao (2003), neural networks Rossi and Conan-Guez (2005); Rossi et al. (2005),  $k$ -nearest neighbors Biau et al. (2005); Laloë (2008), Support Vector Machines (SVM), Rossi and Villa (2006), among a

36 very large variety of methods.

37 In previous works, numerous authors have shown that the derivatives  
38 of the functions lead sometimes to better predictive performances than the  
39 functions themselves in inference tasks, as they provide information about  
40 the shape or the regularity of the function. In particular applications such  
41 as spectrometry Ferraty and Vieu (2006); Rossi et al. (2005); Rossi and Villa  
42 (2006), micro-array data Dejean et al. (2007) and handwriting recognition  
43 Williams et al. (2006); Bahlmann and Burkhardt (2004), these characteristics  
44 lead to accurate predictive models. But, on a theoretical point of the view,  
45 limited results about the effect of the use of the derivatives instead of the  
46 original functions are available: Mas and Pumo (2009) studies this problem  
47 for a linear model built on the first derivatives of the functions. In the present  
48 paper, we also focus on the theoretical relevance of this common practice and  
49 extend Mas and Pumo (2009) to nonparametric approaches and incomplete  
50 knowledge.

51 More precisely, we address the problem of the estimation of the condi-  
52 tional expectation  $\mathbb{E}(Y|X)$  of a random variable  $Y$  given a functional random  
53 variable  $X$ .  $Y$  is assumed to be either real valued (leading to a regression  
54 problem) or to take values in  $\{-1, 1\}$  (leading to a binary classification prob-  
55 lem). We target two theoretical difficulties. The first difficulty is the po-  
56 tential information loss induced by using a derivative instead of the original  
57 function: when one replaces  $X$  by its order  $m$  derivative  $X^{(m)}$ , consistent  
58 estimators (such as kernel models Ferraty and Vieu (2006)) guarantee an  
59 asymptotic estimation of  $\mathbb{E}(Y|X^{(m)})$  but cannot be used directly to address  
60 the original problem, namely estimating  $\mathbb{E}(Y|X)$ . This is a simple conse-  
61 quence of the fact that  $X \mapsto X^{(m)}$  is not a one to one mapping. The second  
62 difficulty is induced by sampling: in practice, functions are never observed  
63 exactly but rather, as explained above, sampled on a discrete sampling grid.  
64 As a consequence, one relies on approximate derivatives,  $\widehat{X}_\tau^{(m)}$  (where  $\tau$  de-  
65 notes the sampling grid). This approach induces even more information loss  
66 with respect to the underlying functional variable  $X$ : in general, a consistent  
67 estimator of  $\mathbb{E}(Y|\widehat{X}_\tau^{(m)})$  will not provide a consistent estimation of  $\mathbb{E}(Y|X)$   
68 and the optimal predictive performances for  $Y$  given  $\widehat{X}_\tau^{(m)}$  will be lower than  
69 the optimal predictive performances for  $Y$  given  $X$ .

70 We show in this paper that the use of a smoothing spline based approach  
71 solves both problems. Smoothing splines are used to estimate the functions  
72 from their sampled version in a convergent way. In addition, properties of

73 splines are used to obtain estimates of the derivatives of the functions with no  
 74 induced information loss. Both aspects are implemented as a preprocessing  
 75 step applied to the multivariate observations generated via the sampling grid.  
 76 The preprocessed observations can then be fed into any finite dimensional  
 77 consistent regression estimator or classifier, leading to a consistent estima-  
 78 tor for the original infinite dimensional problem (in real world applications,  
 79 we instantiate the general scheme in the particular case of kernel machines  
 80 Shawe-Taylor and Cristianini (2004)).

81 The remainder of the paper is organized as follows: Section 2 introduces  
 82 the model, the main smoothness assumption and the notations. Section 3  
 83 recalls important properties of spline smoothing. Section 4 presents approx-  
 84 imation results used to build a general consistent classifier or a general con-  
 85 sistent regression estimator in Section 5. Finally, Section 6 illustrates the  
 86 behavior of the proposed method for two real world spectrometric problems.  
 87 The proofs are given at the end of the article.

## 88 2. Setup and notations

### 89 2.1. Consistent classifiers and regression functions

90 We consider a pair of random variables  $(X, Y)$  where  $X$  takes values in  
 91 a functional space  $\mathcal{X}$  and  $Y$  is either a real valued random variable (regres-  
 92 sion case) or a random variable taking values in  $\{-1, 1\}$  (binary classifica-  
 93 tion case). From this, we are given a learning set  $S_n = \{(X_i, Y_i)\}_{i=1}^n$  of  $n$   
 94 independent copies of  $(X, Y)$ . Moreover, the functions  $X_i$  are not entirely  
 95 known but sampled according to a non random sampling grid of finite length,  
 96  $\tau_d = (t_l)_{l=1}^{|\tau_d|}$ : we only observe  $\mathbf{X}_i^{\tau_d} = (X_i(t_1), \dots, X_i(t_{|\tau_d|}))^T$ , a vector of  $\mathbb{R}^{|\tau_d|}$   
 97 and denote  $S_{n, \tau_d}$  the corresponding learning set. Our goal is to construct:

- 98 1. *in the binary classification case*: a classifier,  $\phi_{n, \tau_d}$ , whose misclassifica-  
 99 tion probability

$$L(\phi_{n, \tau_d}) = \mathbb{P}(\phi_{n, \tau_d}(\mathbf{X}^{\tau_d}) \neq Y)$$

100 asymptotically reaches the Bayes risk

$$L^* = \inf_{\phi: \mathcal{X} \rightarrow \{-1, 1\}} \mathbb{P}(\phi(X) \neq Y)$$

101 i.e.,  $\lim_{|\tau_d| \rightarrow +\infty} \lim_{n \rightarrow +\infty} \mathbb{E}(L(\phi_{n, \tau_d})) = L^*$  ;

102 2. *in the regression case*: a regression function,  $\phi_{n,\tau_d}$ , whose  $L^2$  error

$$L(\phi_{n,\tau_d}) = \mathbb{E}([\phi_{n,\tau_d}(\mathbf{X}^{\tau_d}) - Y]^2)$$

103 asymptotically reaches the minimal  $L^2$  error

$$L^* = \inf_{\phi: \mathcal{X} \rightarrow \mathbb{R}} \mathbb{E}([\phi(\mathbf{X}^{\tau_d}) - Y]^2)$$

104 i.e.,  $\lim_{|\tau_d| \rightarrow +\infty} \lim_{n \rightarrow +\infty} L(\phi_{n,\tau_d}) = L^*$ .

105 This definition implicitly requires  $\mathbb{E}(Y^2) < \infty$  and as a consequence,  
 106 corresponds to a  $L^2$  convergence of  $\phi_{n,\tau_d}$  to the conditional expectation  
 107  $\phi^* = \mathbb{E}(Y|X)$ , i.e., to  $\lim_{|\tau_d| \rightarrow +\infty} \lim_{n \rightarrow +\infty} \mathbb{E}([\phi_{n,\tau_d}(\mathbf{X}^{\tau_d}) - \phi^*(X)]^2) =$   
 108 0.

109 Such  $\phi_{n,\tau_d}$  are said to be (*weakly*) *consistent* Devroye et al. (1996); Györfi  
 110 et al. (2002). We have deliberately used the same notations for the (optimal)  
 111 predictive performances in both the binary classification and the regression  
 112 case. We will call  $L^*$  the Bayes risk even in the case of regression. Most of  
 113 the theoretical background of this paper is common to both the regression  
 114 case and the classification case: the distinction between both cases will be  
 115 made only when necessary.

116 As pointed out in the introduction, the main difficulty is to show that  
 117 the performances of a model built on the  $\mathbf{X}_i^{\tau_d}$  asymptotically reach the best  
 118 performance achievable on the original functions  $X_i$ . In addition, we will  
 119 build the model on derivatives estimated from the  $\mathbf{X}_i^{\tau_d}$ .

## 120 2.2. Smoothness assumption

121 Our goal is to leverage the functional nature of the data by allow-  
 122 ing differentiation operators to be applied to functions prior their submis-  
 123 sion to a more common classifier or regression function. Therefore we as-  
 124 sume that the functional space  $\mathcal{X}$  contains only differentiable functions.  
 125 More precisely,  $\mathcal{X}$  is the Sobolev space  $\mathcal{H}^m = \{h \in L^2([0, 1]) \mid \forall j =$   
 126  $1, \dots, m, D^j h \text{ exists in the weak sense, and } D^m h \in L^2([0, 1])\}$ , where  $D^j h$   
 127 is the  $j$ -th derivative of  $h$  (also denoted by  $h^{(j)}$ ) and for an integer  $m \geq 1$ .  
 128 Of course, by a straightforward generalization, any bounded interval can be  
 129 considered instead of  $[0, 1]$ .

130 To estimate the underlying functions  $X_i$  and their derivatives from sam-  
 131 pled data, we rely on smoothing splines. More precisely, let us consider  
 132 a deterministic function  $x \in \mathcal{H}^m$  sampled on the aforementioned grid. A  
 133 smoothing spline estimate of  $x$  is the solution,  $\hat{x}_{\lambda, \tau_d}$ , of

$$\arg \min_{h \in \mathcal{H}^m} \frac{1}{|\tau_d|} \sum_{l=1}^{|\tau_d|} (x(t_l) - h(t_l))^2 + \lambda \int_{[0,1]} (h^{(m)}(t))^2 dt, \quad (1)$$

134 where  $\lambda$  is a regularization parameter that balances interpolation error and  
 135 smoothness (measured by the  $L^2$  norm of the  $m$ -th derivative of the esti-  
 136 mate). The goal is to show that a classifier or a regression function built  
 137 on  $\hat{X}_{\lambda, \tau_d}^{(m)}$  is consistent for the original problem (i.e., the problem defined by  
 138 the pair  $(X, Y)$ ): this means that using  $\hat{X}_{\lambda, \tau_d}^{(m)}$  instead of  $X$  has no dramatic  
 139 consequences on the accuracy of the classifier or of the regression function.  
 140 In other words, asymptotically, no information loss occurs when one replaces  
 141  $X$  by  $\hat{X}_{\lambda, \tau_d}^{(m)}$ .

142 The proof is based on the following steps:

- 143 1. First, we show that building a classifier or a regression function on  
 144  $\hat{X}_{\lambda, \tau_d}^{(m)}$  is approximately equivalent to building a classifier or a regression  
 145 function on  $\mathbf{X}^{\tau_d} = (X(t_l))_{l=1}^{|\tau_d|}$  using a specific metric. This is done by  
 146 leveraging the Reproducing Kernel Hilbert Space (RKHS) structure of  
 147  $\mathcal{H}^m$ . This part serves one main purpose: it provides a solution to work  
 148 with estimation of the derivatives of the original function in a way  
 149 that preserves all the information available in  $\mathbf{X}^{\tau_d}$ . In other words, the  
 150 best predictive performances for  $Y$  theoretically available by building a  
 151 multivariate model on  $\mathbf{X}^{\tau_d}$  are equal to the best predictive performances  
 152 obtained by building a functional model on  $\hat{X}_{\lambda, \tau_d}^{(m)}$ .
- 153 2. Then, we link  $\mathbb{E}(Y|\hat{X}_{\lambda, \tau_d}^{(m)})$  with  $\mathbb{E}(Y|X)$  by approximation results  
 154 available for smoothing splines. This part of the proof handles the  
 155 effects of sampling.
- 156 3. Finally, we glue both results via standard  $\mathbb{R}^{|\tau_d|}$  consistency results.

157 **3. Smoothing splines and differentiation operators**

158 *3.1. RKHS and smoothing splines*

159 As we want to work on derivatives of functions from  $\mathcal{H}^m$ , a natural inner  
 160 product for two functions of  $\mathcal{H}^m$  would be  $(u, v) \rightarrow \int_0^1 u^{(m)}(t)v^{(m)}(t)dt$ .  
 161 However, we prefer to use an inner product of  $\mathcal{H}^m$  ( $\int_0^1 u^{(m)}(t)v^{(m)}(t)dt$  only  
 162 induces a semi-norm on  $\mathcal{H}^m$ ) because, as will be shown later, such an inner  
 163 product is related to an inner product between the sampled functions  
 164 considered as vectors of  $\mathbb{R}^{|\tau_d|}$ .

165 This can be done by decomposing  $\mathcal{H}^m$  into  $\mathcal{H}^m = \mathcal{H}_0^m \oplus \mathcal{H}_1^m$  Kimeldorf  
 166 and Wahba (1971), where  $\mathcal{H}_0^m = \text{Ker}D^m = \mathbb{P}^{m-1}$  (the space of polynomial  
 167 functions of degree less or equal to  $m - 1$ ) and  $\mathcal{H}_1^m$  is an infinite dimensional  
 168 subspace of  $\mathcal{H}^m$  defined via  $m$  boundary conditions. The boundary condi-  
 169 tions are given by a full rank linear operator from  $\mathcal{H}^m$  to  $\mathbb{R}^m$ , denoted  $B$ , such  
 170 that  $\text{Ker}B \cap \mathbb{P}^{m-1} = \{0\}$ . Classical examples of boundary conditions include  
 171 the case of “natural splines” (for  $m = 2$ ,  $h(0) = h(1) = 0$ ) and constraints  
 172 that target only the first values of  $h$  and its derivatives at a fixed position, for  
 173 instance the conditions:  $h(0) = \dots = h^{(m-1)}(0) = 0$ . Other boundary condi-  
 174 tions can be used Berlinet and Thomas-Agnan (2004); Besse and Ramsay  
 175 (1986); Craven and Wahba (1978), depending on the application.

176 Once the boundary conditions are fixed, an inner product on both  $\mathcal{H}_0^m$   
 177 and  $\mathcal{H}_1^m$  can be defined:

$$\langle u, v \rangle_1 = \langle D^m u, D^m v \rangle_{L^2} = \int_0^1 u^{(m)}(t)v^{(m)}(t)dt$$

178 is an inner product on  $\mathcal{H}_1^m$  (as  $h \in \mathcal{H}_1^m$  and  $D^m h \equiv 0$  give  $h \equiv 0$ ). Moreover,  
 179 if we denote  $B = (B^j)_{j=1}^m$ , then  $\langle u, v \rangle_0 = \sum_{j=1}^m B^j u B^j v$  is an inner product  
 180 on  $\mathcal{H}_0^m$ . We obtain this way an inner product on  $\mathcal{H}^m$  given by

$$\begin{aligned} \langle u, v \rangle_{\mathcal{H}^m} &= \int_0^1 u^{(m)}(t)v^{(m)}(t)dt + \sum_{j=1}^m B^j u B^j v \\ &= \langle \mathcal{P}_1^m(u), \mathcal{P}_1^m(v) \rangle_1 + \langle \mathcal{P}_0^m(u), \mathcal{P}_0^m(v) \rangle_0 \end{aligned}$$

181 where  $\mathcal{P}_i^m$  is the projector on  $\mathcal{H}_i^m$ .

182 Equipped with  $\langle \cdot, \cdot \rangle_{\mathcal{H}^m}$ ,  $\mathcal{H}^m$  is a Reproducing Kernel Hilbert Space  
 183 (RKHS, see e.g. Berlinet and Thomas-Agnan (2004); Heckman and Ramsay  
 184 (2000); Wahba (1990)). More precisely, it exists a kernel  $k : [0, 1]^2 \rightarrow \mathbb{R}$  such

185 that, for all  $u \in \mathcal{H}^m$  and all  $t \in [0, 1]$ ,  $\langle u, k(t, \cdot) \rangle_{\mathcal{H}^m} = u(t)$ . The same occurs  
 186 for  $\mathcal{H}_0^m$  and  $\mathcal{H}_1^m$  which respectively have reproducing kernels denoted by  $k_0$   
 187 and  $k_1$ . We have  $k = k_0 + k_1$ .

188 In the most common cases,  $k_0$  and  $k_1$  have already been explicitly cal-  
 189 culated (see e.g., Berlinet and Thomas-Agnan (2004), especially chapter 6,  
 190 sections 1.1 and 1.6.2). For example, for  $m \geq 1$  and the boundary conditions  
 191  $h(0) = h'(0) = \dots = h^{(m-1)}(0) = 0$ , we have:

$$k_0(s, t) = \sum_{k=0}^{m-1} \frac{t^k s^k}{(k!)^2}.$$

192 and

$$k_1(s, t) = \int_0^1 \frac{(t-w)_+^{m-1} (s-w)_+^{m-1}}{(m-1)!^2} dw.$$

### 193 3.2. Computing the splines

194 We need now to compute to  $\hat{x}_{\lambda, \tau_d}$  starting with  $\mathbf{x}^{\tau_d} = (x(t))_{t \in \tau_d}^T$ . This  
 195 can be done via a theorem from Kimeldorf and Wahba (1971). We need the  
 196 following compatibility assumptions between the sampling grid  $\tau_d$  and the  
 197 boundary conditions operator  $B$ :

198 **Assumption 1.** *The sampling grid  $\tau_d = (t_l)_{l=1}^{|\tau_d|}$  is such that*

- 199 1. *sampling points are distinct in  $[0, 1]$  and  $|\tau_d| \geq m - 1$*
- 200 2. *the  $m$  boundary conditions  $B^j$  are linearly independent from the  $|\tau_d|$*   
 201 *linear forms  $h \mapsto h(t_l)$ , for  $l = 1, \dots, |\tau_d|$  (defined on  $\mathcal{H}^m$ )*

202 Then  $\hat{x}_{\lambda, \tau_d}$  and  $\mathbf{x}^{\tau_d} = (x(t))_{t \in \tau_d}^T$  are linked by the following result:

203 **Theorem 1** (Kimeldorf and Wahba (1971)). *Under Assumption (A1), the*  
 204 *unique solution  $\hat{x}_{\lambda, \tau_d}$  to equation (1) is given by:*

$$\hat{x}_{\lambda, \tau_d} = \mathcal{S}_{\lambda, \tau_d} \mathbf{x}^{\tau_d}, \quad (2)$$

205 where  $\mathcal{S}_{\lambda, \tau_d}$  is a full rank linear operator from  $\mathbb{R}^{|\tau_d|}$  to  $\mathcal{H}^m$  defined by:

$$\mathcal{S}_{\lambda, \tau_d} = \omega^T M_0 + \eta^T M_1 \quad (3)$$

206 with



- 207 •  $M_0 = (U(K_1 + \lambda I_d)^{-1} U^T)^{-1} U(K_1 + \lambda I_d)^{-1}$
- 208 •  $M_1 = (K_1 + \lambda I_d)^{-1} (I_d - U^T M_0)$ ;
- 209 •  $\{\omega_1, \dots, \omega_m\}$  is a basis of  $\mathbb{P}^{m-1}$ ,  $\omega = (\omega_1, \dots, \omega_m)^T$  and  $U =$   
210  $(\omega_i(t))_{i=1, \dots, m}^T_{t \in \tau_d}$ ;
- 211 •  $\eta = (k_1(t, \cdot))_{t \in \tau_d}^T$  and  $K_1 = (k_1(t, t'))_{t, t' \in \tau_d}$ .

### 212 3.3. No information loss

213 The first important consequence of Theorem 1 is that building a model  
214 on  $\widehat{X}_{\lambda, \tau_d}$  or on  $\mathbf{X}^{\tau_d}$  leads to the same optimal predictive performances (to the  
215 same Bayes risk). This is formalized by the following corollary:

216 **Corollary 1.** *Under Assumption (A1), we have*

- 217 • *in the binary classification case:*

$$\inf_{\phi: \mathcal{H}^m \rightarrow \{-1, 1\}} \mathbb{P} \left( \phi(\widehat{X}_{\lambda, \tau_d}) \neq Y \right) = \inf_{\phi: \mathbb{R}^{|\tau_d|} \rightarrow \{-1, 1\}} \mathbb{P} \left( \phi(\mathbf{X}^{\tau_d}) \neq Y \right) \quad (4)$$

- 218 • *in the regression case:*

$$\inf_{\phi: \mathcal{H}^m \rightarrow \mathbb{R}} \mathbb{E} \left( \left[ \phi \left( \widehat{X}_{\lambda, \tau_d} \right) - Y \right]^2 \right) = \inf_{\phi: \mathbb{R}^{|\tau_d|} \rightarrow \mathbb{R}} \mathbb{E} \left( \left[ \phi \left( \mathbf{X}^{\tau_d} \right) - Y \right]^2 \right) \quad (5)$$

### 219 3.4. Differentiation operator

220 The second important consequence of Theorem 1 is that the inner product  
221  $\langle \cdot, \cdot \rangle_{\mathcal{H}^m}$  is equivalent to a specific inner product on  $\mathbb{R}^{|\tau_d|}$  given in the following  
222 corollary:

223 **Corollary 2.** *Under Assumption (A1) and for any  $\mathbf{u}^{\tau_d} = (u(t))_{t \in \tau_d}^T$  and*  
224  $\mathbf{v}^{\tau_d} = (v(t))_{t \in \tau_d}^T$  *in  $\mathbb{R}^{|\tau_d|}$ ,*

$$\langle \widehat{u}_{\lambda, \tau_d}, \widehat{v}_{\lambda, \tau_d} \rangle_{\mathcal{H}^m} = (\mathbf{u}^{\tau_d})^T \mathbf{M}_{\lambda, \tau_d} \mathbf{v}^{\tau_d} \quad (6)$$

225 *where  $\mathbf{M}_{\lambda, \tau_d} = M_0^T W M_0 + M_1^T K_1 M_1$  with  $W = (\langle w_i, w_j \rangle_0)_{i, j=1, \dots, m}$ . The*  
226 *matrix  $\mathbf{M}_{\lambda, \tau_d}$  is symmetric and positive definite and defines an inner product*  
227 *on  $\mathbb{R}^{|\tau_d|}$ .*

228 The corollary is a direct consequence of equations (2) and (3).

229 In practice, the corollary means that the euclidean space  $(\mathbb{R}^{|\tau_d|}, \langle \cdot, \cdot \rangle_{\mathbf{M}_{\lambda, \tau_d}})$   
 230 is isomorphic to  $(\mathcal{I}_{\lambda, \tau_d}, \langle \cdot, \cdot \rangle_{\mathcal{H}^m})$ , where  $\mathcal{I}_{\lambda, \tau_d}$  is the image of  $\mathbb{R}^{|\tau_d|}$  by  $\mathcal{S}_{\lambda, \tau_d}$ . As  
 231 a consequence, one can use the Hilbert structure of  $\mathcal{H}^m$  directly in  $\mathbb{R}^{|\tau_d|}$  via  
 232  $\mathbf{M}_{\lambda, \tau_d}$ : as the inner product of  $\mathcal{H}^m$  is defined on the order  $m$  derivatives of the  
 233 functions, this corresponds to using those derivatives instead of the original  
 234 functions.

235 More precisely, let  $\mathbf{Q}_{\lambda, \tau_d}$  be the transpose of the Cholesky triangle of  $\mathbf{M}_{\lambda, \tau_d}$   
 236 (given by the Cholesky decomposition  $\mathbf{Q}_{\lambda, \tau_d}^T \mathbf{Q}_{\lambda, \tau_d} = \mathbf{M}_{\lambda, \tau_d}$ ). Corollary 2  
 237 shows that  $\mathbf{Q}_{\lambda, \tau_d}$  acts as an approximate differentiation operation on sampled  
 238 functions.

239 Let us indeed consider an estimation method for multivariate inputs based  
 240 only on inner products or norms (that are directly derived from the inner  
 241 products), such as, e.g., Kernel Ridge Regression Saunders et al. (1998);  
 242 Shawe-Taylor and Cristianini (2004). In this latter case, if a Gaussian kernel  
 243 is used, the regression function has the following form:

$$u \mapsto \sum_{i=1}^n T_i \alpha_i e^{-\gamma \|U_i - u\|_{\mathbb{R}^p}^2} \quad (7)$$

244 where  $(U_i, T_i)_{1 \leq i \leq n}$  are learning examples in  $\mathbb{R}^p \times \{-1, 1\}$  and the  $\alpha_i$  are non  
 245 negative real values obtained by solving a quadratic programming problem  
 246 and  $\gamma$  is a parameter of the method. Then, if we use Kernel Ridge Regression  
 247 on the training set  $\{(\mathbf{Q}_{\lambda, \tau_d} \mathbf{X}_i^{\tau_d}, Y_i)\}_{i=1}^n$  (rather than the original training set  
 248  $\{(\mathbf{X}_i^{\tau_d}, Y_i)\}_{i=1}^n$ ), it will work on the norm in  $L^2$  of the derivatives of order  
 249  $m$  of the spline estimates of the  $X_i$  (up to the boundary conditions). More  
 250 precisely, the regression function will have the following form:

$$\begin{aligned} \mathbf{x}^{\tau_d} &\mapsto \sum_{i=1}^n Y_i \alpha_i e^{-\gamma \|\mathbf{Q}_{\lambda, \tau_d} \mathbf{X}_i^{\tau_d} - \mathbf{Q}_{\lambda, \tau_d} \mathbf{x}^{\tau_d}\|_{\mathbb{R}^{|\tau_d|}}^2} \\ &\mapsto \sum_{i=1}^n Y_i \alpha_i e^{-\gamma \|D^m \widehat{X}_{i, \lambda, \tau_d} - D^m \widehat{x}_{\lambda, \tau_d}\|_{L^2}^2} \\ &\quad \times e^{-\gamma \sum_{j=1}^m (B^j \widehat{X}_{i, \lambda, \tau_d} - B^j \widehat{x}_{\lambda, \tau_d})^2} \end{aligned}$$

251 In other words, up to the boundary conditions, an estimation method based  
 252 solely on inner products, or on norms derived from these inner products,

253 can be given modified inputs that will make it work on an estimation of the  
 254 derivatives of the observed functions.

255 **Remark 1.** As shown in Corollary 1 in the previous section, building a  
 256 model on  $\mathbf{X}^{\tau_d}$  or on  $\widehat{X}_{\lambda, \tau_d}$  leads to the same optimal predictive performances.  
 257 In addition, it is obvious that given any one-to-one mapping  $f$  from  $\mathbb{R}^{|\tau_d|}$  to  
 258 itself, building a model on  $f(\mathbf{X}^{\tau_d})$  gives also the same optimal performances  
 259 than building a model on  $\mathbf{X}^{\tau_d}$ . Then as  $\mathbf{Q}_{\lambda, \tau_d}$  is invertible, the optimal  
 260 predictive performances achievable with  $\mathbf{Q}_{\lambda, \tau_d} \mathbf{X}^{\tau_d}$  are equal to the optimal  
 261 performances achievable with  $\mathbf{X}^{\tau_d}$  or with  $\widehat{X}_{\lambda, \tau_d}$ .

262 In practice however, the actual preprocessing of the data can have a strong  
 263 influence on the obtained performances, as will be illustrated in Section 6.  
 264 The goal of the theoretical analysis of the present section is to guarantee  
 265 that no systematic loss can be observed as a consequence of the proposed  
 266 functional preprocessing scheme.

#### 267 4. Approximation results

268 The previous section showed that working on  $\mathbf{X}^{\tau_d}$ ,  $\mathbf{Q}_{\lambda, \tau_d} \mathbf{X}^{\tau_d}$  or  $\widehat{X}_{\lambda, \tau_d}$   
 269 makes no difference in terms of optimal predictive performances. The present  
 270 section addresses the effects of sampling: asymptotically, the optimal predic-  
 271 tive performances obtained on  $\widehat{X}_{\lambda, \tau_d}$  converge to the optimal performances  
 272 achievable on the original and unobserved functional variable  $X$ .

##### 273 4.1. Spline approximation

274 From the sampled random function  $\mathbf{X}^{\tau_d} = (X(t_1), \dots, X(t_{|\tau_d|}))$ , we can  
 275 build an estimate,  $\widehat{X}_{\lambda, \tau_d}$ , of  $X$ . To ensure consistency, we must guarantee  
 276 that  $\widehat{X}_{\lambda, \tau_d}$  converges to  $X$ . In the case of a deterministic function  $x$ , this  
 277 problem has been studied in numerous papers, such as Craven and Wahba  
 278 (1978); Ragozin (1983); Cox (1984); Utreras (1988); Wahba (1990) (among  
 279 others). Here we recall one of the results which is particularly well adapted  
 280 to our context.

281 Obviously, the sampling grid must behave correctly, whereas the infor-  
 282 mation contained in  $\mathbf{X}^{\tau_d}$  will not be sufficient to recover  $X$ . We need also  
 283 the regularization parameter  $\lambda$  to depend on  $\tau_d$ . Following Ragozin (1983),  
 284 a sampling grid  $\tau_d$  is characterized by two quantities:

$$\begin{aligned} \overline{\Delta}_{\tau_d} &= \max\{t_1, t_2 - t_1, \dots, 1 - t_{|\tau_d|}\} \\ \underline{\Delta}_{\tau_d} &= \min_{1 \leq i < |\tau_d|} \{t_{i+1} - t_i\}. \end{aligned} \tag{8}$$

285 One way to control the distance between  $X$  and  $\widehat{X}_{\lambda, \tau_d}$  is to bound the ratio  
 286  $\overline{\Delta}_{\tau_d} / \underline{\Delta}_{\tau_d}$  so as to ensure quasi-uniformity of the sampling grid.

287 More precisely, we will use the following assumption:

288 **Assumption 2.** *There is  $R$  such that  $\overline{\Delta}_{\tau_d} / \underline{\Delta}_{\tau_d} \leq R$  for all  $d$ .*

289 Then we have:

290 **Theorem 2** (Ragozin (1983)). *Under Assumptions (A1) and (A2), there are*  
 291 *two constants  $A_{R,m}$  and  $B_{R,m}$  depending only on  $R$  and  $m$ , such that for any*  
 292  *$x \in \mathcal{H}^m$  and any positive  $\lambda$ :*

$$\|\widehat{x}_{\lambda, \tau_d} - x\|_{L^2}^2 \leq \left( A_{R,m} \lambda + B_{R,m} \frac{1}{|\tau_d|^{2m}} \right) \|D^m x\|_{L^2}^2.$$

293 This result is a rephrasing of Corollary 4.16 from Ragozin (1983) which  
 294 is itself a direct consequence of Theorem 4.10 from the same paper.

295 Convergence of  $\widehat{x}_{\lambda, \tau_d}$  to  $x$  is then obtained by the following simple as-  
 296 sumptions:

297 **Assumption 3.** *The series of sampling points  $\tau_d$  and the series of regular-*  
 298 *ization parameters,  $\lambda$ , depending on  $\tau_d$  and denoted by  $(\lambda_d)_{d \geq 1}$ , are such that*  
 299  *$\lim_{d \rightarrow +\infty} |\tau_d| = +\infty$  and  $\lim_{d \rightarrow +\infty} \lambda_d = 0$ .*

#### 300 4.2. Conditional expectation approximation

301 The next step consists in relating the optimal predictive performances  
 302 for the regression and the classification problem  $(X, Y)$  to the performances  
 303 associated to  $(\widehat{X}_{\lambda_d, \tau_d}, Y)$  when  $d$  goes to infinity, i.e., relating  $L^*$  to

304 1. *binary classification case:*

$$L_d^* = \inf_{\phi: \mathcal{H}^m \rightarrow \{-1, 1\}} \mathbb{P} \left( \phi(\widehat{X}_{\lambda_d, \tau_d}) \neq Y \right),$$

305 2. *regression case:*

$$L_d^* = \inf_{\phi: \mathcal{H}^m \rightarrow \mathbb{R}} \mathbb{E} \left( [\phi(\widehat{X}_{\lambda_d, \tau_d}) - Y]^2 \right)$$

306 Two sets of assumptions will be investigated to provide the convergence  
 307 of the Bayes risk  $L_d^*$  to  $L^*$ :

308 **Assumption 4. Either**

309 (A4a)  $\mathbb{E}(\|D^m X\|_{L^2}^2)$  is finite and  $Y \in \{-1, 1\}$ ,

310 **or**

311 (A4b)  $\tau_d \subset \tau_{d+1}$  and  $\mathbb{E}(Y^2)$  is finite.

312 The first assumption (A4a) requires an additional smoothing property for  
 313 the predictor functional variable  $X$  and is only valid for a binary classifica-  
 314 tion problem whereas the second assumption (A4b) requires an additional  
 315 property for the sampling point series: they have to be growing sets.

316 Theorem 2 then leads to the following corollary:

317 **Corollary 3.** *Under Assumptions (A1)-(A4), we have:*

$$\lim_{d \rightarrow +\infty} L_d^* = L^*.$$

## 318 5. General consistent functional classifiers and regression functions

### 319 5.1. Definition of classifiers and regression functions on derivatives

320 Let us now consider any consistent classification or regression scheme for  
 321 standard multivariate data based either on the inner product or on the Eu-  
 322 clidean distance between observations. Examples of such classifiers are Sup-  
 323 port Vector Machine Steinwart (2002), the kernel classification rule Devroye  
 324 and Krzyżak (1989) and  $k$ -nearest neighbors Devroye and Györfi (1985);  
 325 Zhao (1987) to name a few. In the same way, multilayer perceptrons Lu-  
 326 gosi and Zeger (1990), kernel estimates Devroye and Krzyżak (1989) and  
 327  $k$ -nearest neighbors regression Devroye et al. (1994) are consistent regression  
 328 estimators. Additional examples of consistent estimators in classification and  
 329 regression can be found in Devroye et al. (1996); Györfi et al. (2002).

330 We denote  $\psi_{\mathcal{D}}$  the estimator constructed by the chosen scheme using a  
 331 dataset  $\mathcal{D} = \{(U_i, T_i)_{1 \leq i \leq n}\}$ , where the  $(U_i, T_i)_{1 \leq i \leq n}$  are  $n$  independent copies  
 332 of a pair of random variables  $(U, T)$  with values in  $\mathbb{R}^p \times \{-1, 1\}$  (classification)  
 333 or  $\mathbb{R}^p \times \mathbb{R}$  (regression).

334 The proposed functional scheme consists in choosing the estimator  $\phi_{n, \tau_d}$   
 335 as  $\psi_{\mathcal{E}_{n, \tau_d}}$  with the dataset  $\mathcal{E}_{n, \tau_d}$  defined by:

$$\mathcal{E}_{n, \tau_d} = \{(\mathbf{Q}_{\lambda_d, \tau_d} \mathbf{X}_i^{\tau_d}, Y_i)_{1 \leq i \leq n}\}$$

336 As pointed out in Section 3.4, the linear transformation  $\mathbf{Q}_{\lambda_d, \tau_d}$  is an approx-  
 337 imate multivariate differentiation operator: up to the boundary conditions,  
 338 an estimator based on  $\mathbf{Q}_{\lambda_d, \tau_d} \mathbf{X}^{\tau_d}$  is working on the  $m$ -th derivative of  $\widehat{X}_{\lambda_d, \tau_d}$ .

339 In more algorithmic terms, the estimator is obtained as follows:

- 340 1. choose an appropriate value for  $\lambda_d$
- 341 2. compute  $\mathbf{M}_{\lambda_d, \tau_d}$  using Theorem 1 and Corollary 2;
- 342 3. compute the Cholesky decomposition of  $\mathbf{M}_{\lambda_d, \tau_d}$  and the transpose of  
 343 the Cholesky triangle,  $\mathbf{Q}_{\lambda_d, \tau_d}$  (such that  $\mathbf{Q}_{\lambda_d, \tau_d}^T \mathbf{Q}_{\lambda_d, \tau_d} = \mathbf{M}_{\lambda_d, \tau_d}$ );
- 344 4. compute  $\mathbf{Q}_{\lambda_d, \tau_d} \mathbf{X}_i^{\tau_d}$  to obtain the transformed dataset  $\mathcal{E}_{n, \tau_d}$ ;
- 345 5. build a classifier/regression function  $\psi_{\mathcal{E}_{n, \tau_d}}$  with a multivariate method  
 346 in  $\mathbb{R}^{|\tau_d|}$  applied to the dataset  $\mathcal{E}_{n, \tau_d}$ ;
- 347 6. associate to a new sampled function  $\mathbf{X}_{n+1}^{\tau_d}$  the prediction  
 348  $\psi_{\mathcal{E}_{n, \tau_d}}(\mathbf{Q}_{\lambda, \tau_d} \mathbf{X}_{n+1}^{\tau_d})$ .

349 Figure 5.1 illustrates the way the method performs: instead of relying  
 350 on an approximation of the function and then on the derivation preprocess-  
 351 ing of this estimates, it directly uses an equivalent metric by applying the  
 352  $\mathbf{Q}_{\lambda_d, \tau_d}$  matrix to the sampled function. The consistency result proved in The-  
 353 orem 3 shows that, combined with any consistent multidimensional learning  
 354 algorithm, this method is (asymptotically) equivalent to using the original  
 355 function drawn at the top left side of Figure 5.1.

356 On a practical point of view, Wahba (1990) demonstrates that cross val-  
 357 idated estimates of  $\lambda$  achieve suitable convergence rates. Hence, steps 1 and  
 358 2 can be computed simultaneously by minimizing the total cross validated  
 359 error for all the observations, given by

$$\sum_{i=1}^n \frac{1}{|\tau_d|} \sum_{t \in \tau_d} \frac{(x_i(t) - \widehat{x}_{i, \lambda, \tau_d}(t))^2}{(1 - A_{tt}(\lambda))^2},$$

360 where  $A$  is a  $|\tau_d| \times |\tau_d|$  matrix called the *influence matrix* (see Wahba (1990)),  
 361 over a finite number of  $\lambda$  values.




Figure 1: Method scheme and its equivalence to the usual approach for using derivatives in learning algorithms.

362 *5.2. Consistency result*

363 Corollary 1 and Corollary 3 guarantee that the estimator proposed in the  
364 previous section is consistent:

365 **Theorem 3.** *Under assumptions (A1)-(A4), the series of classi-*  
366 *fiers/regression functions  $(\phi_{n,\tau_d})_{n,d}$  is consistent:*

$$\lim_{d \rightarrow +\infty} \lim_{n \rightarrow +\infty} \mathbb{E} (L\phi_{n,\tau_d}) = L^*$$

367 *5.3. Discussion*

368 While Theorem 3 is very general, it could be easily extended to cover  
369 special cases such as additional hypothesis needed by the estimation scheme  
370 or to provide data based parameter selections. We discuss briefly those issues  
371 in the present section.

372 It should first be noted that most estimation schemes,  $\psi_{\mathcal{D}}$ , depend on  
373 parameters that should fulfill some assumptions for the scheme to be con-  
374 sistent. For instance, in the Kernel Ridge Regression method in  $\mathbb{R}^p$ , with  
375 Gaussian kernel,  $\psi_{\mathcal{D}}$  has the form given in Equation (7) where the  $(\alpha_i)$  are  
376 the solutions of

$$\arg \min_{\alpha \in \mathbb{R}^n} \sum_{i=1}^n \left( T_i - \sum_{j=1}^n T_j \alpha_j e^{-\gamma \|U_i - U_j\|_{\mathbb{R}^p}^2} \right)^2 +$$

$$\delta_n \sum_{i,j=1}^n T_i T_j \alpha_i \alpha_j e^{-\gamma \|U_i - U_j\|_{\mathbb{R}^p}^2}.$$

377 The method thus depends on the parameter of the Gaussian kernel,  $\gamma$  and  
378 of the regularization parameter  $\delta_n$ . This method is known to be consistent if  
379 (see Theorem 9.1 of Steinwart and Christmann (2008)):

$$\delta_n \xrightarrow{n \rightarrow +\infty} 0 \quad \text{and} \quad n\delta_n^4 \xrightarrow{n \rightarrow +\infty} +\infty.$$

380 Additional conditions of this form can obviously be directly integrated in  
381 Theorem 3 to obtain consistency results specific to the corresponding algo-  
382 rithms.

383 Moreover, practitioners generally rely on data based selection of the pa-  
384 rameters of the estimation scheme  $\psi_{\mathcal{D}}$  via a validation method: for instance,  
385 rather than setting  $\delta_n$  to e.g.,  $n^{-5}$  for  $n$  observations (a choice which is com-  
386 patible with theoretical constraints on  $\delta_n$ ), one chooses the value of  $\delta_n$  that



387 optimizes an estimation of the performances of the regression function ob-  
388 tained on an independent data set (or via a re-sampling approach).

389 In addition to the parameters of the estimation scheme, functional data  
390 raise the question of the convenient order of the derivative,  $m$ , and of the  
391 sampling grid optimality. In practical applications, the number of available  
392 sampling points can be unnecessarily large (see Biau et al. (2005) for an ex-  
393 ample with more than 8 000 sampling points). The preprocessing performed  
394 by  $\mathbf{Q}_{\lambda_d, \tau_d}$  do not change the dimensionality of the data which means that  
395 overfitting can be observed in practice when the number of sampling points  
396 is large compared to the number of functions. Moreover, processing very  
397 high dimensional vectors is time consuming. It is there quite interesting in  
398 practice to use a down-sampled version of the original grid.

399 To select the parameters of  $\psi_D$ , the order of the derivative and/or the  
400 down-sampled grid, a validation strategy, based on splitting the dataset into  
401 training and validation sets, could be used. A simple adaptation of the idea  
402 of Berline et al. (2008); Biau et al. (2005); Laloë (2008); Rossi and Villa  
403 (2006) shows that a penalized validation method can be used to choose any  
404 combination of those parameters consistently. According to those papers,  
405 the condition for the consistency of the validation strategy would simply  
406 relate the shatter coefficients of the set of classifiers in  $\mathbb{R}^d$  to the penalization  
407 parameter of the validation. Once again, this type of results is a rather direct  
408 extension of Theorem 3.

## 409 6. Applications

410 In this section, we show that the proposed approach works as expected on  
411 real world spectrometric examples: for some applications, the use of deriva-  
412 tives leads to more accurate models than the direct processing of the spectra  
413 (see e.g. Rossi et al. (2005); Rossi and Villa (2006) for other examples of such  
414 a behavior based on ad hoc estimators of the spectra derivatives). It should  
415 be noted that the purpose of this section is only to illustrate the behavior  
416 of the proposed method on finite datasets. The theoretical results of the  
417 present paper show that all consistent schemes have asymptotically identical  
418 performances, and therefore that using derivatives is asymptotically useless.  
419 On a finite dataset however, preprocessing can have strong influence on the  
420 predictive performances, as will be illustrated in the present section. In ad-  
421 dition, schemes that are not universally consistent, e.g., linear models, can  
422 lead to excellent predictive performances on finite datasets; such models are

423 therefore included in the present section despite the fact the theory does not  
424 apply to them.

### 425 *6.1. Methodology*

426 The methodology followed for the two illustrative datasets is roughly the  
427 same:

- 428 1. the dataset is randomly split into a training set on which the model is  
429 estimated and a test set on which performances are computed. The split  
430 is repeated several times. The Tecator dataset (Section 6.2) is rather  
431 small (240 spectra) and exhibits a rather large variability in predic-  
432 tive performances between different random splits. We have therefore  
433 used 250 random splits. For the Yellow-berry dataset (Section 6.3), we  
434 used only 50 splits as the relative variability in performances is far less  
435 important.
- 436 2.  $\lambda$  is chosen by a global leave-one-out strategy on the spectra contained  
437 in training set (as suggested in Section 5.1). More precisely, a leave-one-  
438 out estimate of the reconstruction error of the spline approximation of  
439 each training spectrum is computed for a finite set of candidate values  
440 for  $\lambda$ . Then a common  $\lambda$  is chosen by minimizing the average over  
441 the training spectra of the leave-one-out reconstruction errors. This  
442 choice is relevant as cross validation estimates of  $\lambda$  are known to have  
443 favorable theoretical properties (see Craven and Wahba (1978); Utreras  
444 (1981) among others).
- 445 3. for regression problems, a Kernel Ridge Regression (KRR) Saunders  
446 et al. (1998); Shawe-Taylor and Cristianini (2004) is then performed to  
447 estimate the regression function; this method is consistent when used  
448 with a Gaussian kernel under additional conditions on the parameters  
449 (see Theorem 9.1 of Steinwart and Christmann (2008)); as already ex-  
450 plained, in the applications, Kernel Ridge Regression is performed both  
451 with a Gaussian kernel and with a linear kernel (in that last case, the  
452 model is essentially a ridge regression model). Parameters of the models  
453 (a regularization parameter,  $\delta_n$ , in all cases and a kernel parameter,  $\gamma$   
454 for Gaussian kernels) are chosen by a grid search that minimizes a vali-  
455 dation based estimate of the performances of the model (on the training  
456 set). A leave-one-out solution has been chosen: in Kernel Ridge Re-  
457 gression, the leave-one-out estimate of the performances of the model is

458 obtained as a by-product of the estimation process, without additional  
459 computation cost, see e.g. Cawley and Talbot (2004).

460 Additionally, for a sake of comparison with a more traditional approach  
461 in FDA, Kernel Ridge Regression is compared with a nonparametric  
462 kernel estimate for the Tecator dataset (Section 6.2.1). Nonparametric  
463 kernel estimate is the first nonparametric approach introduced in Func-  
464 tional Data Analysis Ferraty and Vieu (2006) and can thus be seen as  
465 a basis for comparison in the context of regression with functional pre-  
466 dictors. For this method, the same methodology as with Kernel Ridge  
467 Regression was used: the parameter of the model (i.e., the bandwidth)  
468 was selected on a grid search minimizing a cross-validation estimate of  
469 the performances of the model. In this case, a 4-fold cross validation  
470 estimate was used instead of a leave-one-out estimate to avoid a large  
471 computational cost.

472 4. for the classification problem, a Support Vector Machine (SVM) is used  
473 Shawe-Taylor and Cristianini (2004). As KRR, SVM are consistent  
474 when used with a Gaussian kernel Steinwart (2002). We also use a  
475 SVM with a linear kernel as this is quite adapted for classification in  
476 high dimensional spaces associated to sampled function data. We also  
477 use a K-nearest neighbor model (KNN) for reference. Parameters of the  
478 models (a regularization parameter for both SVM, a kernel parameter,  
479  $\gamma$  for Gaussian kernels and number of neighbors  $K$  for KNN) are chosen  
480 by a grid search that minimizes a validation based estimate of the  
481 classification error: we use a 4-fold cross-validation to get this estimate.

482 5. We evaluate the models obtained for each random split on the test set.  
483 We report the mean and the standard deviation of the performance  
484 index (classification error and mean squared error, respectively) and  
485 assess the significance of differences between the reported figures via  
486 paired Student tests (with level 1%).

487 6. Finally, we compare models estimated on the raw spectra and on spec-  
488 tra transformed via the  $\mathbf{Q}_{\lambda_d, \tau_d}$  matrix for  $m = 1$  (first derivative) and  
489  $m = 2$  (second derivative). For both values of  $m$ , we used the most  
490 classical boundary conditions ( $x(0) = 0$  and  $Dx(0) = 0$ ). Depending of  
491 the problem, other boundary conditions could be investigated but this  
492 is outside the scope of the present paper (see Besse and Ramsay (1986);  
493 Heckman and Ramsay (2000) for discussion on this subject). For the

494 Tecator problem, we also compare these approaches with models es-  
495 timated on first and second derivatives based on interpolating splines  
496 (i.e. with  $\lambda = 0$ ) and on first and second derivatives estimated by finite  
497 differences.

498 Note that the kind of preprocessing used has almost no impact on  
499 the computation time. In general, selecting the parameters of the  
500 model with leave-one-out or cross-validation will use significantly more  
501 computing power than constructing the splines and calculating their  
502 derivatives. For instance, computing the optimal  $\lambda$  with the approach  
503 described above takes less than 0.1 second for the Tecator dataset on a  
504 standard PC using our R implementation which is negligible compared  
505 to the several minutes used to select the optimal parameters of the  
506 models used on the preprocessed data.

## 507 6.2. Tecator dataset

508 The first studied dataset is the standard Tecator dataset Thodberg (1996)  
509 <sup>1</sup>. It consists in spectrometric data from the food industry. Each of the  
510 240 observations is the near infrared absorbance spectrum of a meat sample  
511 recorded on a Tecator Infratec Food and Feed Analyzer. Each spectrum is  
512 sampled at 100 wavelengths uniformly spaced in the range 850–1050 nm.  
513 The composition of each meat sample is determined by analytic chemistry  
514 and percentages of moisture, fat and protein are associated this way to each  
515 spectrum.

516 The Tecator dataset is a widely used benchmark in Functional Data Anal-  
517 ysis, hence the motivation for its use for illustrative purposes. More precisely,  
518 in Section 6.2.1, we address the original regression problem by predicting the  
519 percentage of fat content from the spectra with various regression method  
520 and various estimates of the derivative preprocessing: this analysis shows  
521 that both the method and the use of derivative have a strong effect on the  
522 performances whereas the way the derivatives are estimated has almost no  
523 effect. Additionally, in Section 6.2.2, we apply a noise (with various vari-  
524 ances) to the original spectra in order to study the influence of smoothing  
525 in the case of noisy predictors: this section shows the relevance of the use of  
526 a smoothing spline approach when the data are noisy. Finally, Section 6.2.3  
527 deals with a classification problem derived from the original Tecator problem

---

<sup>1</sup>Data are available on statlib at <http://lib.stat.cmu.edu/datasets/tecator>

528 (in the same way as what was done in Ferraty and Vieu (2003)): conclusions  
529 of this section are similar to the ones of the regression study.

### 530 6.2.1. Fat content prediction

531 As explained above, we first address the regression problem that consists  
532 in predicting the fat content of peaces of meat from the Tecator dataset. The  
533 parameters of the model are optimized with a grid search using the leave-one-  
534 out estimate of the predictive performances (both models use a regularization  
535 parameter, with an additional width parameter in the Gaussian kernel case).  
536 The original data set is split randomly into 160 spectra for learning and 80  
537 spectra for testing. As shown in the result Table 1, the data exhibit a rather  
538 large variability; we use therefore 250 random split to assess the differences  
539 between the different approaches.

540 The performance indexes are the mean squared error (M.S.E.) and the  
541  $R^2$ .<sup>2</sup> As a reference, the target variable (fat) has a variance equal to 14.36.  
542 Results are summarized in Table 1.

543 The first conclusion is that the method itself has a strong effect on the  
544 performances of the prediction: for this application, a linear method is not  
545 appropriate (mean squared errors are much greater for linear methods than  
546 for the kernel ridge regression used with a Gaussian kernel) and the non-  
547 parametric kernel estimate gives worse performances than the kernel ridge  
548 regression (indeed, they are about 10 times worse). Nevertheless, for non-  
549 parametric approaches (Gaussian KKR and NKE), the use of derivatives  
550 has also a strong impact on the performances: for kernel ridge regression,  
551 e.g., preprocessing by estimating the first order derivative leads to a strong  
552 decrease of the mean squared error.

553 Differences between the average MSEs are not always significant, but  
554 we can nevertheless rank the methods in increasing order of modeling error  
555 (using notations explained in Table 1) for Gaussian kernel ridge regression:

$$\text{FD1} \leq \text{IS1} \leq \text{S1} < \text{DF2} \leq \text{SS2} < \text{IS2} < \text{O}$$

556 where  $<$  corresponds to a significant difference (for a paired Student test  
557 with level 1%) and  $\leq$  to a non significant one. In this case, the data are very  
558 smooth and thus the use of smoothing splines instead of a finite differences

---

<sup>2</sup> $R^2 = 1 - \frac{\text{M.S.E.}}{\text{Var}(y)}$  where  $\text{Var}(y)$  is the (empirical) variance of the target variable on the test set.

Method	Data	Average M.S.E. and SD	Average $R^2$
KRR Linear	O	8.69 (4.47)	95.7%
	S1	8.09 (3.85)	96.1%
	IS1	8.09 (3.85)	96.1%
	FD1	8.27 (4.17)	96.0%
	S2	9.64 (4.98)	95.3%
	IS2	9.87 (5.84)	95.2%
	FD2	8.45 (4.18)	95.9%
KRR Gaussian	O	5.02 (11.47)	97.6%
	S1	0.485 (0.385)	99.8%
	IS1	0.485 (0.385)	99.8%
	FD1	<b>0.484</b> (0.387)	<b>99.8%</b>
	S2	0.584 (0.303)	99.7%
	IS2	0.586 (0.303)	99.7%
	FD2	0.569 (0.281)	99.7%
NKE	O	73.1 (16.5)	64.2%
	S1	4.59 (1.09)	97.7%
	IS1	4.59 (1.09)	97.7%
	FD1	4.59 (1.09)	97.7%
	S2	3.75 (1.22)	98.2%
	IS2	3.75 (1.22)	98.2%
	FD2	3.67 (1.18)	98.2%

Table 1: Summary of the performances of the chosen models on the test set (fat Tecator regression problem) when using either a kernel ridge regression (KRR) with linear kernel or with Gaussian kernel or when using a nonparametric kernel estimate (NKE) with various inputs: O (original data), S1 (smoothing splines with order 1 derivatives), IS1 (interpolating splines with order 1 derivatives), FD1 (order 1 derivatives estimated by finite differences) and S2, IS2 and FD2 (the same as previously with order 2 derivatives).

559 approximation does not have a significant impact on the predictions. How-  
 560 ever, in this case, the roughest approach, consisting in the estimation of the  
 561 derivatives by finite differences, gives the best performances.

### 562 6.2.2. Noisy spectra

563 This section studies the situation in which functional data observations  
 564 are corrupted by noise. This is done by adding a noise to each spectrum of  
 565 the Tecator dataset. More precisely, each spectrum has been corrupted by

$$X_i^b(t) = X_i(t) + \epsilon_{it} \quad (9)$$

566 where  $(\epsilon_{it})$  are i.i.d. Gaussian variables with standard deviation equal to  
 567 either 0.01 (small noise) or to 0.2 (large noise). 10 observations of the data  
 568 generated this way are given in Figure 2.

569 The same methodology as for the non noisy data has been applied to  $(X_i^b)$   
 570 to predict the fat content. The experiments have been restricted to the use of  
 571 kernel ridge regression with a Gaussian kernel (according to the nonlinearity  
 572 of the problem shown in the previous section). Results are summarized in  
 573 Table 2 and Figure 3.

574 In addition, the results can be ranked this way:

#### **Noise with sd equal to 0.01**

$$S2 < S1 < IS1 \leq O < FD1 < IS2 \leq FD2$$

#### **Noise with sd equal to 0.2**

$$S1 < O < S2 < FD1 < IS1 < IS2 \leq FD2$$

575 where  $<$  corresponds to a significant difference (for a paired Student test  
 576 with level 1%).

577 The first conclusion of these experiments is that, even though the deriva-  
 578 tives are the relevant predictors, their performances are strongly affected by  
 579 the noise (compared to the ones of the original data: note that the average  
 580 M.S.E. reported in Table 1 are more 10 times lower than the best ones from  
 581 Table 2 and that, in the best cases,  $R^2$  is slightly greater than 50% for the  
 582 most noisy dataset). In particular, using interpolating splines or finite differ-  
 583 ence derivatives leads to highly deteriorated performances. In this situation,  
 584 the approach proposed in the paper is particularly useful and helps to keep




Figure 2: 10 observations of the noisy data generated from the Tecator spectra as in Equation 9



Noise	Data	Average M.S.E. and SD	Average $R^2$
sd = 0.01	O	13.3 (13.5)	93.5%
	S1	7.45 (1.5)	96.4%
	IS1	12.72 (2.2)	93.8%
	FD1	20.03 (2.8)	90.3%
	S2	<b>6.83</b> (1.4)	<b>96.7%</b>
	IS2	31.23 (5.9)	84.9%
	FD2	31.10 (5.9)	84.9%
sd = 0.2	O	87.9 (13.9)	57.4%
	S1	<b>85.0</b> (12.5)	<b>58.8%</b>
	IS1	210.1 (36.1)	-1.9%
	FD1	209.1 (33.0)	-1.4%
	S2	95.9 (12.8)	53.5%
	IS2	213.7 (33.1)	-3.6%
	FD2	235.1 (222.7)	-14.0%

Table 2: Summary of the performances of the chosen models on the test set (fat Tecator regression problem) with noisy spectra.




Figure 3: Mean squared errors boxplot for the noisy fat Tecator regression problem with Gaussian kernel (the worst test samples for  $S$  and  $FD$  have been removed for a sake of clarity)

585 better performances than with the original data. Indeed, the differences of  
586 the smoothing splines approach with the original data is still significant (for  
587 both derivatives in the “small noise” case and for the first order derivative  
588 in the “high noise” case), even though, the most noisy the data are, the  
589 most difficult it is to estimate the derivatives in an accurate way. That is,  
590 except for smoothing spline derivatives, the estimation of the derivatives for  
591 the most noisy dataset is so bad that it leads to negative  $R^2$  when used in  
592 the regression task.

### 593 *6.2.3. Fat content classification*

594 In this section, the fat content regression problem is transformed into a  
595 classification problem. To avoid imbalance in class sizes, the median value  
596 of the fat in the dataset is used as the splitting criterion: the first class  
597 consists in 119 samples with strictly less than 13.5 % of fat, while the second  
598 class contains the other 121 samples with a fat content equal or higher than  
599 13.5 %.

600 As in previous sections, the analysis is conducted on 250 random splits of  
601 the dataset into 160 learning spectra and 80 test spectra. We used stratified  
602 sampling: the test set contains 40 examples from each class. The 4 fold  
603 cross-validation used to select the parameters of the models on the learning  
604 set is also stratified with roughly 20 examples of each class in each fold.

605 The performance index is the mis-classification rate (MCR) on the test  
606 set, reported in percentage and averaged over the 250 random splits. Results  
607 are summarized in Table 3. As in the previous sections, both the model  
608 and the preprocessing have some influence on the results. In particular,  
609 using derivatives always improves the classification accuracy while the actual  
610 method used to compute those derivatives has no particular influence on the  
611 results. Additionally, using interpolation splines leads, in this particular  
612 problem, to results that are exactly identical to the ones obtained with the  
613 smoothing splines: they are not reported in Table 3.

614 More precisely, for the three models (linear SVM, Gaussian SVM and  
615 KNN), differences in mis-classification rates between the smoothing spline  
616 preprocessing and the finite differences calculation is never significant, ac-  
617 cording to a Student test with level 1 %. Additionally while the actual aver-  
618 age mis-classification rates might seem quite different, the large variability of  
619 the results (shown by the standard deviations) leads to significant differences  
620 only for the most obvious cases. In particular, SVM models using derivatives  
621 (of order one or two) are indistinguishable one from another using a Student

Method	Data	Average MCR	SD of MCR
Linear SVM	O	1.41	1.55
	S1	<b>0.73</b>	1.15
	FD1	0.74	1.15
	S2	0.94	1.27
	FD2	0.92	1.23
Gaussian SVM	O	3.39	2.57
	S1	0.97	1.41
	FD1	0.98	1.42
	S2	0.99	2.00
	FD2	0.97	1.27
KNN	O	22.0	5.02
	S1	6.67	2.55
	FD1	6.57	2.55
	S2	1.93	1.65
	FD2	1.93	1.63

Table 3: Summary of the performances of the chosen models on the test set (Tecator fat classification problem). See Table 1 for notations. MCR stands for mis-classification rate, SD for standard deviation.

622 test with level 1 %: all methods with less than 1 % of mean mis-classification  
623 rate perform essentially identically. Other differences are significant: for in-  
624 stance the linear SVM used on raw data performs significantly worse than  
625 any SVM model used on derivatives.

626 It should be noted that the classification task studied in the present sec-  
627 tion is obviously simpler than the regression task from which it is derived.  
628 This explains the very good predictive performances obtained by simple mod-  
629 els such as a linear SVM, especially with the proper preprocessing.

### 630 6.3. Yellow-berry dataset

631 The goal of the last experiment is to predict the presence of yellow-berry in  
632 durum wheat (*Triticum durum*) kernels via a near infrared spectral analysis  
633 (see Figure 4). Yellow-berry is a defect of the durum wheat seeds that reduces  
634 the quality of the flour produced from affected wheat. The traditional way  
635 to assess the occurrence of yellow-berry is by visual analysis of a sample of  
636 the seed stock. In the current application, a quality measure related to the  
637 occurrence of yellow-berry is predicted from the spectrum of the seed.




Figure 4: 20 observations of NIR spectra of durum wheat

638 The dataset consists in 953 spectra sampled at 1049 wavelengths uni-  
639 formly spaced in the range 400–2498 nm. The dataset is split randomly into  
640 600 learning spectra and 353 test spectra. Comparatively to the Tecator  
641 dataset, the variability of the results is smaller in the present case. We used  
642 therefore 50 random splits rather than 250 in the previous section.

643 The regression models were build via a Kernel Ridge Regression approach  
644 using a linear kernel and a Gaussian kernel. In both cases, the regularization  
645 parameter of the model is optimized by a leave-one-out approach. In addi-  
646 tion, the width parameter of the Gaussian kernel is optimized via the same  
647 procedure at the same time.

648 The performance index is the mean squared error (M.S.E.). As a refer-  
649 ence, the target variable has a variance of 0.508. Results are summarized in  
Table 4 and Figure 5.

Kernel and Data	Average M.S.E.	Standard deviation	Average $R^2$
Linear-O	0.122	$8.77 \cdot 10^{-3}$	76.1%
Linear-S1	0.138	$9.53 \cdot 10^{-3}$	73.0%
Linear-S2	0.122	$8.41 \cdot 10^{-3}$	76.1%
Gaussian-O	0.110	$20.2 \cdot 10^{-3}$	78.5%
Gaussian-S1	0.0978	$7.92 \cdot 10^{-3}$	80.9%
Gaussian-S2	0.0944	$8.35 \cdot 10^{-3}$	81.5%

Table 4: Summary of the performances of the chosen models on the test set (durum wheat regression problem)

650 As in the previous section, we can rank the methods in increasing order  
651 of modelling error, we obtain the following result:  
652

$$\text{G-S2} < \text{G-S1} < \text{G-O} < \text{L-O} \leq \text{L-S2} < \text{L-S1},$$

653 where G stands for Gaussian kernel and L for linear kernel (hence G-S2 stands  
654 for kernel ridge regression with gaussian kernel and smoothing splines with  
655 order 2 derivatives);  $<$  corresponds to a significant difference (for a paired  
656 Student test with level 1%) and  $\leq$  to a non significant one. For this appli-  
657 cation, there is a significant gain in using a non linear model (the Gaussian  
658 kernel). In addition, the use of derivatives leads to less contrasted perfor-  
659 mances that the ones obtained in the previous section but it still improves  
660 the quality of the non linear model in a significant way. In term of normal-  
661 ized mean squared error (mean squared error divided by the variance of the




Figure 5: Mean squared error boxplots for the “durum wheat” regression problem (see Table 4 for the full names of the regression models)

662 target variable), using a non linear model with the second derivatives of the  
663 spectra corresponds to an average gain of more than 5% (i.e., a reduction of  
664 the normalised mean squared error from 24% for the standard linear model  
665 to 18.6%).

## 666 7. Conclusion

667 In this paper we proposed a theoretical analysis of a common practice that  
668 consists in using derivatives in classification or regression problems when the  
669 predictors are curves. Our method relies on smoothing splines reconstruction  
670 of the functions which are known only via a discrete deterministic sampling.  
671 The method is proved to be consistent for very general classifiers or regres-  
672 sion schemes: it reaches asymptotically the best risk that could have been  
673 obtained by constructing a regression/classification model on the true ran-  
674 dom functions.

675 We have validated the approach by combining it with nonparametric re-  
676 gression and classification algorithms to study two real-world spectrometric  
677 datasets. The results obtained in these applications confirm once again that  
678 relying on derivatives can improve the quality of predictive models compared  
679 to a direct use of the sampled functions. The way the derivatives are esti-  
680 mated does not have a strong impact on the performances except when the  
681 data are noisy. In this case, the use of smoothing splines is quite relevant.

682 In the future, several issues could be addressed. An important practical  
683 problem is the choice of the best order of the derivative,  $m$ . We consider  
684 that a model selection approach relying on a penalized error loss could be  
685 used, as is done, in e.g., Rossi and Villa (2006), to select the dimension of  
686 truncated basis representation for functional data. Note that in practice,  
687 such parameter selection method could lead to select  $m = 0$  and therefore to  
688 automatically exclude derivative calculation when it is not needed. This will  
689 extend the application range of the proposed model.

690 A second important point to study is the convergence rate for the method.  
691 It would be very convenient for instance, to be able to relate the size of  
692 the sampling grid to the number of functions. But, this latter issue would  
693 require the use of additional assumptions on the smoothness of the regression  
694 function whereas the result presented in this paper, even if more limited, only  
695 needs mild conditions.



696 **8. Acknowledgement**

697 We thank Cécile Levasseur and Sylvain Coulomb (École d'Ingénieurs de  
698 Purpan, EIP, Toulouse, France) for sharing the interesting problem presented  
699 in Section 6.3.

700 We also thank Philippe Besse (Institut de Mathématiques de Toulouse,  
701 Université de Toulouse, France) for helpful discussions and suggestions.

702 Finally, we thank the anonymous reviewers for their valuable comments  
703 and suggestions that helped to improve the quality of the paper.

704 **References**

705 Bahlmann, C., Burkhardt, H., 2004. The writer independent online hand-  
706 writing recognition system *frog on hand* and cluster generative statistical  
707 dynamic time warping. *IEEE Transactions on Pattern Analysis and Ma-*  
708 *chine Intelligence* 26, 299–310.

709 Berline, A., Biau, G., Rouvière, L., 2008. Functional supervised classifica-  
710 tion with wavelets. *Annales de l'ISUP* 52, 61–80.

711 Berline, A., Thomas-Agnan, C., 2004. *Reproducing Kernel Hilbert Spaces*  
712 *in Probability and Statistics*. Kluwer Academic Publisher.

713 Besse, P., Ramsay, J., 1986. Principal component analysis of sampled curves.  
714 *Psychometrika* 51, 285–311.

715 Biau, G., Bunea, F., Wegkamp, M., 2005. Functional classification in Hilbert  
716 spaces. *IEEE Transactions on Information Theory* 51, 2163–2172.

717 Cardot, H., Ferraty, F., Sarda, P., 1999. Functional linear model. *Statistics*  
718 *and Probability Letters* 45, 11–22.

719 Cawley, G., Talbot, N., 2004. Fast exact leave-one-out cross-validation of  
720 sparse least-squares support vector machines. *Neural Networks* 17, 1467–  
721 1475.

722 Cox, D., 1984. Multivariate smoothing splines functions. *SIAM Journal on*  
723 *Numerical Analysis* 21, 789–813.

724 Craven, P., Wahba, G., 1978. Smoothing noisy data with spline functions.  
725 *Numerische Mathematik* 31, 377–403.

- 726 Dauxois, J., Pousse, A., 1976. Les analyses factorielles en calcul des proba-  
727 bilités et en statistique : essai d'étude synthétique. Thèse d'État. Univer-  
728 sité Toulouse III.
- 729 Dejean, S., Martin, P., Baccini, A., Besse, P., 2007. Clustering time-series  
730 gene expression data using smoothing spline derivatives. EURASIP Jour-  
731 nal on Bioinformatics and Systems Biology 2007, Article ID70561.
- 732 Deville, J., 1974. Méthodes statistiques et numériques de l'analyse har-  
733 monique. Annales de l'INSEE 15, 3–97.
- 734 Devroye, L., Györfi, L., 1985. Nonparametric Density Estimation: the  $L_1$   
735 view. John Wiley, New York.
- 736 Devroye, L., Györfi, L., Krzyżak, A., Lugosi, G., 1994. On the strong uni-  
737 versal consistancy of nearest neighbor regression function estimates. The  
738 Annals of Statistics 22, 1371–1385.
- 739 Devroye, L., Györfi, L., Lugosi, G., 1996. A Probabilistic Theory for Pattern  
740 Recognition. Springer-Verlag, New York.
- 741 Devroye, L., Krzyżak, A., 1989. An equivalence theorem for  $l_1$  convergence  
742 of the kernel regression estimate. Journal of Statistical Planning and In-  
743 ference 23, 71–82.
- 744 Faragó, T., Györfi, L., 1975. On the continuity of the error distortion func-  
745 tion for multiple-hypothesis decisions. IEEE Transactions on Information  
746 Theory 21, 458–460.
- 747 Ferraty, F., Vieu, P., 2003. Curves discrimination: a non parametric ap-  
748 proach. Computational and Statistical Data Analysis 44, 161–173.
- 749 Ferraty, F., Vieu, P., 2006. NonParametric Functional Data Analysis.  
750 Springer.
- 751 Ferré, L., Yao, A., 2003. Functional sliced inverse regression analysis. Statis-  
752 tics 37, 475–488.
- 753 Györfi, L., Kohler, M., Krzyżak, A., Walk, H., 2002. A Distribution-Free  
754 Theory of Nonparametric Regression. Springer, New York.

- 755 Heckman, N., Ramsay, J., 2000. Penalized regression with model-based  
756 penalties. *The Canadian Journal of Statistics* 28, 241–258.
- 757 James, G., 2002. Generalized linear models with functional predictor vari-  
758 ables. *Journal of the Royal Statistical Society Series B* 64, 411–432.
- 759 James, G., Hastie, T., 2001. Functional linear discriminant analysis for ir-  
760 regularly sampled curves. *Journal of the Royal Statistical Society, Series*  
761 *B* 63, 533–550.
- 762 James, G., Hastie, T., Sugar, C., 2000. Principal component models for  
763 sparse functional data. *Biometrika* 87, 587–602.
- 764 James, G., Silverman, B., 2005. Functional adaptive model estimation. *Jour-*  
765 *nal of the American Statistical Association* 100, 565–576.
- 766 Kallenberg, O., 1997. *Foundations of Modern Probability. Probability and*  
767 *its Applications*, Springer.
- 768 Kimeldorf, G., Wahba, G., 1971. Some results on Tchebycheffian spline  
769 functions. *Journal of Mathematical Analysis and Applications* 33, 82–95.
- 770 Laloë, T., 2008. A k-nearest neighbor approach for functional regression.  
771 *Statistics and Probability Letters* 78, 1189–1193.
- 772 Lugosi, G., Zeger, K., 1990. Nonparametric estimation via empirical risk  
773 minimization. *IEEE Transaction on Information Theory* 41, 677–687.
- 774 Mas, A., Pumo, B., 2009. Functional linear regression with derivatives. *Jour-*  
775 *nal of Nonparametric Statistics* 21, 19–40. Submitted: under revision.  
776 Available at <http://www.math.univ-montp2.fr/~mas/FLRD.pdf>.
- 777 Pollard, D., 2002. *A User’s Guide to Measure Theoretic Probability*. Cam-  
778 *bridge University Press*, Cambridge.
- 779 Ragozin, D., 1983. Error bounds for derivative estimation based on spline  
780 smoothing of exact or noisy data. *Journal of Approximation Theory* 37,  
781 335–355.
- 782 Ramsay, J., Dalzell, C., 1991. Some tools for functional data analysis (with  
783 discussion). *Journal of the Royal Statistical Society. Series B. Statistical*  
784 *Methodology* 53, 539–572.

- 785 Ramsay, J., Silverman, B., 1997. *Functional Data Analysis*. Springer Verlag,  
786 New York.
- 787 Ramsay, J., Silverman, B., 2002. *Applied Functional Data Analysis*. Springer  
788 Verlag.
- 789 Rossi, F., Conan-Guez, B., 2005. Functional multi-layer perceptron: a non-  
790 linear tool for functional data analysis. *Neural Networks* 18, 45–60.
- 791 Rossi, F., Conan-Guez, B., 2006. Theoretical properties of projection based  
792 multilayer perceptrons with functional inputs. *Neural Processing Letters*  
793 23, 55–70.
- 794 Rossi, F., Delannay, N., Conan-Guez, B., Verleysen, M., 2005. Representa-  
795 tion of functional data in neural networks. *Neurocomputing* 64, 183–210.
- 796 Rossi, F., Villa, N., 2006. Support vector machine for functional data classi-  
797 fication. *Neurocomputing* 69, 730–742.
- 798 Saunders, G., Gammerman, A., Vovk, V., 1998. Ridge regression learning  
799 algorithm in dual variables, in: *Proceedings of the Fifteenth International*  
800 *Conference on Machine Learning (ICML'98)*, Madison, Wisconsin, USA.  
801 pp. 515–521.
- 802 Shawe-Taylor, J., Cristianini, N., 2004. *Kernel methods for pattern analysis*.  
803 Cambridge University Press, Cambridge, UK.
- 804 Steinwart, I., 2002. Support vector machines are universally consistent. *Jour-  
805 nal of Complexity* 18, 768–791.
- 806 Steinwart, I., Christmann, A., 2008. *Support Vector Machines*. Information  
807 Science and Statistics, Springer.
- 808 Thodberg, H., 1996. A review of bayesian neural network with an application  
809 to near infrared spectroscopy. *IEEE Transaction on Neural Networks* 7,  
810 56–72.
- 811 Utreras, F., 1981. Optimal smoothing of noisy data using spline functions.  
812 *SIAM Journal on Scientific Computing* 2, 153–163.
- 813 Utreras, F., 1988. Boundary effects on convergence rates for Tikhonov regu-  
814 larization. *Journal of Approximation Theory* 54, 235–249.

- 815 Wahba, G., 1990. Spline Models for Observational Data. Society for Indus-  
816 trial and Applied Mathematics, Philadelphia, Pennsylvania.
- 817 Williams, B., Toussaint, M., Storkey, A., 2006. Extracting motion primitives  
818 from natural handwriting data, in: In Proceedings of the International  
819 Conference on Artificial Neural Networks (ICANN).
- 820 Zhao, L., 1987. Exponential bounds of mean error for the nearest neighbor  
821 estimates of regression functions. *Journal of Multivariate Analysis* 21,  
822 168–178.

## 823 9. Proofs

### 824 9.1. Theorem 1

825 In the original theorem (Lemma 3.1) in Kimeldorf and Wahba (1971),  
826 one has to verify that  $(k_0(t_l, \cdot))_l$  spans  $\mathcal{H}_0^m$  and that  $(k_1(t_l, \cdot))_l$  are linearly  
827 independent. These are consequences of Assumption (A1).

828 First,  $k_0(s, t) = \sum_{i,j=0}^{m-1} b_{ij}^{(-1)} s^i t^j$  where  $\tilde{B} = (b_{i,j}^{(-1)})_{i,j}$  is the in-  
829 verse of  $(\sum_{l=1}^m B^l s^i B^l t^j)_{i,j}$  (see Heckman and Ramsay (2000)). Then  
830  $(k_0(t_1, s), \dots, k_0(t_{|\tau_d|}, s)) = (1, s, \dots, s^{m-1}) \tilde{B} [V_{m-1}(t_1, \dots, t_{|\tau_d|})]^T$  where  
831  $V_{m-1}(t_1, \dots, t_{|\tau_d|})$  is the Vandermonde matrix with  $m - 1$  columns and  $|\tau_d|$   
832 rows associated to values  $t_1, \dots, t_{|\tau_d|}$ . If the  $(t_l)_l$  are distinct, this matrix is  
833 of full rank.

834 Moreover the reproducing property shows that  $\sum_{l=1}^{|\tau_d|} a_l k_1(t_l, \cdot) \equiv 0$  im-  
835 plies  $\sum_{l=1}^{|\tau_d|} a_l f(t_l) \equiv 0$  for all  $f \in \mathcal{H}_1^m$ . Hence,  $\mathcal{H}_1^m = \text{Ker}(B^T, \sum_{l=1}^{\tau_d} a_l \zeta_l)^T$   
836 where  $\zeta_l$  denotes the linear form  $h \in \mathcal{H}^m \rightarrow h(t_l)$ . As the co-dimension of  
837  $\mathcal{H}_1^m$  is  $\dim \mathcal{H}_0^m = m$  and as, by Assumption (A1),  $B$  is linearly independent  
838 of  $\sum_{l=1}^{\tau_d} a_l \zeta_l$ , we thus have  $\sum_{l=1}^{\tau_d} a_l \zeta_l \equiv 0$  (or  $\text{codim Ker}(B^T, \sum_{l=1}^{\tau_d} a_l \zeta_l)^T =$   
839  $\dim \text{Im}(B^T, \sum_{l=1}^{\tau_d} a_l \zeta_l)$  would be  $m+1$ ). Thus, we obtain that  $\sum_{l=1}^{|\tau_d|} a_l f(t_l) \equiv$   
840  $0$  for all  $f$  in  $\mathcal{H}^m$  and, as  $(t_l)$  are distinct, that  $a_l = 0$  for all  $l$ , leading to the  
841 independence conclusion for the  $(k_1(t_l, \cdot))_l$ .

842 Finally, we prove that  $\mathcal{S}_{\lambda, \tau_d}$  is of full rank. Indeed, if  $\mathcal{S}_{\lambda, \tau_d} \mathbf{x}^{\tau_d} = 0$ ,  
843  $\omega^T M_0 \mathbf{x}^{\tau_d} = 0$  and  $\eta^T M_1 \mathbf{x}^{\tau_d} = 0$ . As  $(\omega_k)_k$  is a basis of  $\mathcal{H}_0^m$ ,  $\omega^T M_0 \mathbf{x}^{\tau_d} = 0$   
844 implies  $M_0 \mathbf{x}^{\tau_d} = 0$  and therefore  $M_1 = (K_1 + \lambda I_d)^{-1}$ . As shown above,  
845 the  $(k_1(t_l, \cdot))_l$  are linearly independent and therefore  $\eta M_1 \mathbf{x}^{\tau_d} = 0$  implies  
846  $M_1 \mathbf{x}^{\tau_d} = 0$ , which in turns leads to  $\mathbf{x}^{\tau_d} = 0$  via the simplified formula for  $M_1$ .

847 *9.2. Corollary 1*

848 We give only the proof for the classification case, the regression case is  
849 identical.

850 According to Theorem 1, there is a full rank linear mapping from  $\mathbb{R}^{|\tau_d|}$   
851 to  $\mathcal{H}^m$ ,  $\mathcal{S}_{\lambda, \tau_d}$ , such that for any function  $x \in \mathcal{H}^m$ ,  $\hat{x}_{\lambda, \tau_d} = \mathcal{S}_{\lambda, \tau_d} \mathbf{x}^{\tau_d}$ . Let  
852 us denote  $\mathcal{I}_{\lambda, \tau_d}$  the image of  $\mathbb{R}^{|\tau_d|}$  by  $\mathcal{S}_{\lambda, \tau_d}$ ,  $\mathbf{P}_{\lambda, \tau_d}$  the orthogonal projection  
853 from  $\mathcal{H}^m$  to  $\mathcal{I}_{\lambda, \tau_d}$  and  $\mathcal{S}_{\lambda, \tau_d}^{-1}$  the inverse of  $\mathcal{S}_{\lambda, \tau_d}$  on  $\mathcal{I}_{\lambda, \tau_d}$ . Obviously, we have  
854  $\mathcal{S}_{\lambda, \tau_d}^{-1} \circ \mathbf{P}_{\lambda, \tau_d}(\hat{x}_{\lambda, \tau_d}) = \mathbf{x}^{\tau_d}$ .

855 Let  $\psi$  be a measurable function from  $\mathbb{R}^{|\tau_d|}$  to  $\{-1, 1\}$ . Then  $\zeta_\psi$  de-  
856 fined on  $\mathcal{H}^m$  by  $\zeta_\psi(u) = \psi(\mathcal{S}_{\lambda, \tau_d}^{-1} \circ \mathbf{P}_{\lambda, \tau_d}(u))$  is a measurable function from  
857  $\mathcal{H}^m$  to  $\{-1, 1\}$  (because  $\mathcal{S}_{\lambda, \tau_d}^{-1}$  and  $\mathbf{P}_{\lambda, \tau_d}$  are both continuous). Then for  
858 any measurable  $\psi$ ,  $\inf_{\phi: \mathcal{H}^m \rightarrow \{-1, 1\}} \mathbb{P}(\phi(\hat{X}_{\lambda, \tau_d}) \neq Y) \leq \mathbb{P}(\zeta_\psi(\hat{X}_{\lambda, \tau_d}) \neq Y) =$   
859  $\mathbb{P}(\psi(\mathbf{X}^{\tau_d}) \neq Y)$ , and therefore

$$\inf_{\phi: \mathcal{H}^m \rightarrow \{-1, 1\}} \mathbb{P}(\phi(\hat{X}_{\lambda, \tau_d}) \neq Y) \leq \inf_{\phi: \mathbb{R}^{|\tau_d|} \rightarrow \{-1, 1\}} \mathbb{P}(\phi(\mathbf{X}^{\tau_d}) \neq Y). \quad (10)$$

860 Conversely, let  $\psi$  be a measurable function from  $\mathcal{H}^m$  to  $\{-1, 1\}$ . Then  $\zeta_\psi$  de-  
861 fined on  $\mathbb{R}^{|\tau_d|}$  by  $\zeta_\psi(\mathbf{u}) = \psi(\mathcal{S}_{\lambda, \tau_d}(\mathbf{u}))$ , is measurable. Then for any measurable  
862  $\psi$ ,  $\inf_{\phi: \mathbb{R}^{|\tau_d|} \rightarrow \{-1, 1\}} \mathbb{P}(\phi(\mathbf{X}^{\tau_d}) \neq Y) \leq \mathbb{P}(\zeta_\psi(\mathbf{X}^{\tau_d}) \neq Y) = \mathbb{P}(\psi(\hat{X}_{\lambda, \tau_d}) \neq Y)$ ,  
863 and therefore

$$\inf_{\phi: \mathbb{R}^{|\tau_d|} \rightarrow \{-1, 1\}} \mathbb{P}(\phi(\mathbf{X}^{\tau_d}) \neq Y) \leq \inf_{\phi: \mathcal{H}^m \rightarrow \{-1, 1\}} \mathbb{P}(\phi(\hat{X}_{\lambda, \tau_d}) \neq Y). \quad (11)$$

864 The combination of equations (10) and (11) gives equality (4).

865 *9.3. Corollary 3*

866 **1. Suppose assumption (A4a) is fulfilled**

867 The proof is based on Theorem 1 in Faragó and Györfi (1975). This  
868 theorem relates the Bayes risk of a classification problem based on  
869  $(X, Y)$  with the Bayes risk of the problem  $(T_d(X), Y)$  where  $(T_d)$  is a  
870 series of transformations on  $X$ .

871 More formally, for a pair of random variables  $(X, Y)$ , where  $X$  takes  
872 values in  $\mathcal{X}$ , an arbitrary metric space, and  $Y$  in  $\{-1, 1\}$ , let us

873 denote for any series of functions  $T_d$  from  $\mathcal{X}$  to itself,  $L^*(T_d) =$   
 874  $\inf_{\phi: \mathcal{X} \rightarrow \{-1,1\}} \mathbb{P}(\phi(T_d(X)) \neq Y)$ . Theorem 1 from Faragó and Györfi  
 875 (1975) states that  $\mathbb{E}(\delta(T_d(X), X)) \xrightarrow{d \rightarrow +\infty} 0$  implies  $L^*(T_d) \xrightarrow{d \rightarrow +\infty} L^*$ ,  
 876 where  $\delta$  denotes the metric on  $\mathcal{X}$ .

877 This can be applied to  $\mathcal{X} = (\mathcal{H}^m, \langle \cdot, \cdot \rangle_{L^2})$  with  $T_d(X) =$   
 878  $\widehat{X}_{\lambda_d, \tau_d} = S_{\lambda_d, \tau_d} \mathbf{X}^{\tau_d}$ : under Assumptions (A1) and (A2), Theo-  
 879 rem 2 gives:  $\|T_d(X) - X\|_{L^2}^2 \leq \left( A_{R,m} \lambda_d + B_{R,m} \frac{1}{|\tau_d|^{2m}} \right) \|D^m X\|_{L^2}^2$ .  
 880 Taking the expectation of both sides gives  $\mathbb{E}(\|T_d(X) - X\|_{L^2}^2) \leq$   
 881  $\left( A_{R,m} \lambda_d + B_{R,m} \frac{1}{|\tau_d|^{2m}} \right) \mathbb{E}(\|D^m X\|_{L^2}^2)$ , using the fact that the constants  
 882 are independent of the function under analysis. Then under Assump-  
 883 tions (A4a) and (A3),  $\mathbb{E}(\|T_d(X) - X\|_{L^2}^2) \xrightarrow{d \rightarrow +\infty} 0$ . According to  
 884 Faragó and Györfi (1975), this implies  $\lim_{d \rightarrow \infty} L_d^* = L^*$ .

## 885 2. Suppose assumption (A4b) is fulfilled

886 The conclusion will follow both for classification case and for regression  
 887 case. The proof follows the general ideas of Biau et al. (2005); Rossi  
 888 and Conan-Guez (2006); Rossi and Villa (2006); Laloë (2008). Under  
 889 assumption (A1), by Theorem 1 and with an argument similar to those  
 890 developed in the proof of Corollary 1,  $\sigma(\widehat{X}_{\lambda_d, \tau_d}) = \sigma(\{X(t)\}_{t \in \tau_d})$ . From  
 891 assumption (A4b),  $\sigma(\{X(t)\}_{t \in \tau_d})$  is clearly a filtration. Moreover, as  
 892  $\mathbb{E}(Y)$  and thus  $\mathbb{E}(Y^2)$  are finite,  $\mathbb{E}(Y|\widehat{X}_{\lambda_d, \tau_d})$  is a uniformly bounded  
 893 martingale for this filtration (see Lemma 35 of Pollard (2002)). This  
 894 martingale converges in  $L^1$ -norm to  $\mathbb{E}(Y|\sigma(\cup_d \sigma(\widehat{X}_{\lambda_d, \tau_d})))$ ; we have

- 895 •  $\sigma(\cup_d \sigma(\widehat{X}_{\lambda_d, \tau_d})) \subset \sigma(X)$  as  $\widehat{X}_{\lambda_d, \tau_d}$  is a function of  $X$  (via Theo-  
 896 rem 1);
- 897 • by Theorem 2,  $\widehat{X}_{\lambda_d, \tau_d} \xrightarrow{d \rightarrow +\infty, \text{ surely}} X$  in  $L^2$  which proves that  $X$   
 898 is  $\sigma(\cup_d \sigma(\widehat{X}_{\lambda_d, \tau_d}))$ -measurable.

899 Finally,  $\mathbb{E}(Y|\sigma(\cup_d \sigma(\widehat{X}_{\lambda_d, \tau_d}))) = \mathbb{E}(Y|X)$  and  
 900  $\mathbb{E}(Y|\widehat{X}_{\lambda_d, \tau_d}) \xrightarrow{d \rightarrow +\infty, L^1} \mathbb{E}(Y|X)$ .

901 The conclusion follows from the fact that:

902 (a) *binary classification case*: the bound  $L_d^* - L^* \leq$   
 903  $2\mathbb{E} \left( \left| \mathbb{E} \left( Y | \widehat{X}_{\lambda_d, \tau_d} \right) - \mathbb{E} (Y | X) \right| \right)$  (see Theorem 2.2 of Devroye  
 904 et al. (1996)) concludes the proof;

905 (b) *regression case*: as  $\mathbb{E} (Y^2)$  is finite,  $\mathbb{E} \left( \mathbb{E} \left( Y | \widehat{X}_{\lambda_d, \tau_d} \right)^2 \right)$  is also fi-  
 906 nite and the convergence also happens for the quadratic norm (see  
 907 Corollary 6.22 in Kallenberg (1997)), i.e.,

$$\lim_{d \rightarrow +\infty} \mathbb{E} \left( \left( \mathbb{E} (Y | X) - \mathbb{E} \left( Y | \widehat{X}_{\lambda_d, \tau_d} \right) \right)^2 \right) = 0$$

908 Hence, as  $L_d^* - L^* = \mathbb{E} \left( \left( \mathbb{E} (Y | X) - \mathbb{E} \left( Y | \widehat{X}_{\lambda_d, \tau_d} \right) \right)^2 \right)$ , the con-  
 909 clusion follows.

#### 910 9.4. Theorem 3

911 We have

$$L(\phi_{n,d}) - L^* = L\phi_{n,\tau_d} - L_d^* + L_d^* - L^*. \quad (12)$$

912 Let  $\epsilon$  be a positive real. By Corollary 3, it exists  $d_0 \in \mathbb{N}^*$  such that, for all  
 913  $d \geq d_0$ ,

$$L_d^* - L^* \leq \epsilon. \quad (13)$$

914 Moreover, as shown in Corollary 1 and as  $\mathbf{Q}_{\lambda_d, \tau_d}$  is invertible, we have  
 915 in the binary classification case:  $L_d^* = \inf_{\phi: \mathbb{R}^{|\tau_d|} \rightarrow \{-1,1\}} \mathbb{P}(\phi(\mathbf{X}^{\tau_d}) \neq Y) =$   
 916  $\inf_{\phi: \mathbb{R}^{|\tau_d|} \rightarrow \{-1,1\}} \mathbb{P}(\phi(\mathbf{Q}_{\lambda_d, \tau_d} \mathbf{X}^{\tau_d}) \neq Y)$ , and in the regression case:  $L_d^* =$   
 917  $\inf_{\phi: \mathbb{R}^{|\tau_d|} \rightarrow \mathbb{R}} \mathbb{E}([\phi(\mathbf{X}^{\tau_d}) - Y]^2) = \inf_{\phi: \mathbb{R}^{|\tau_d|} \rightarrow \mathbb{R}} \mathbb{E}([\phi(\mathbf{Q}_{\lambda_d, \tau_d} \mathbf{X}^{\tau_d}) - Y]^2)$ . By hy-  
 918 pothesis, for any fixed  $d$ ,  $\phi_{n,\tau_d}$  is consistent, that is

$$\lim_{n \rightarrow +\infty} \mathbb{E}(L(\phi_{n,\tau_d})) = \inf_{\phi: \mathbb{R}^{|\tau_d|} \rightarrow \{-1,1\}} \mathbb{P}(\phi(\mathbf{Q}_{\lambda_d, \tau_d} \mathbf{X}^{\tau_d}) \neq Y),$$

919 in the classification case and

$$\lim_{n \rightarrow +\infty} \mathbb{E}(L(\phi_{n,\tau_d})) = \inf_{\phi: \mathbb{R}^{|\tau_d|} \rightarrow \mathbb{R}} \mathbb{E}([\phi(\mathbf{Q}_{\lambda_d, \tau_d} \mathbf{X}^{\tau_d}) - Y]^2),$$

920 in the regression case, and therefore for any fixed  $d_0$ ,  
 921  $\lim_{n \rightarrow +\infty} \mathbb{E} \left( L(\phi_{n,\tau_{d_0}}) \right) = L_{d_0}^*$ . Combined with equations (12) and  
 922 (13), this concludes the proof.